



## Estimating fisheries reference points from catch and resilience

Rainer Froese<sup>1</sup>, Nazli Demirel<sup>2</sup>, Gianpaolo Coro<sup>3</sup>, Kristin M Kleisner<sup>4</sup> & Henning Winker<sup>5,6</sup>

<sup>1</sup>GEOMAR Helmholtz Centre for Ocean Research Kiel, Düsternbrooker Weg 20, 24105 Kiel, Germany; <sup>2</sup>Institute of Marine Sciences and Management, Istanbul University, 34134 Istanbul, Turkey; <sup>3</sup>Istituto di Scienza e Tecnologie dell'Informazione "A. Faedo", Consiglio Nazionale delle Ricerche (CNR), via Moruzzi 1, 56124 Pisa, Italy; <sup>4</sup>National Oceanographic and Atmospheric Administration, Northeast Fisheries Science Center, 166 Water St., Woods Hole, MA 02543, USA; <sup>5</sup>South African National Biodiversity Institute, Kirstenbosch Research Centre, 7735 Claremont, South Africa; <sup>6</sup>Centre for Statistics in Ecology, Environment and Conservation (SEEC), Department of Statistical Sciences, University of Cape Town, Private Bag X3, 7700 Rondebosch, South Africa

### Abstract

This study presents a Monte Carlo method (CMSY) for estimating fisheries reference points from catch, resilience and qualitative stock status information on data-limited stocks. It also presents a Bayesian state-space implementation of the Schaefer production model (BSM), fitted to catch and biomass or *catch-per-unit-of-effort* (CPUE) data. Special emphasis was given to derive informative priors for productivity, unexploited stock size, catchability and biomass from population dynamics theory. Both models gave good predictions of the maximum intrinsic rate of population increase  $r$ , unexploited stock size  $k$  and maximum sustainable yield  $MSY$  when validated against simulated data with known parameter values. CMSY provided, in addition, reasonable predictions of relative biomass and exploitation rate. Both models were evaluated against 128 real stocks, where estimates of biomass were available from full stock assessments. BSM estimates of  $r$ ,  $k$  and  $MSY$  were used as benchmarks for the respective CMSY estimates and were not significantly different in 76% of the stocks. A similar test against 28 data-limited stocks, where CPUE instead of biomass was available, showed that BSM and CMSY estimates of  $r$ ,  $k$  and  $MSY$  were not significantly different in 89% of the stocks. Both CMSY and BSM combine the production model with a simple stock–recruitment model, accounting for reduced recruitment at severely depleted stock sizes.

### Correspondence

Rainer Froese,  
GEOMAR Helmholtz  
Centre for Ocean  
Research Kiel,  
Düsternbrooker Weg  
20, 24105 Kiel,  
Germany  
Tel.:  
+494316004579  
Fax:  
+494316001699  
E-mail: rfroese@  
geomar.de

Received 18 Feb  
2016

Accepted 11 Aug  
2016

**Keywords** Bayesian state-space model, biomass dynamic model, data-limited stock assessment, Monte Carlo method, stock–recruitment relationship, surplus production model

<b>Introduction</b>	<b>507</b>
<b>Material and methods</b>	<b>507</b>
General description of method	508
Selection of real stocks and generation of simulated stocks	509
CMSY analysis	510
<i>Determining the boundaries of the r-k space</i>	510

Setting prior biomass ranges	510
Finding viable <i>r-k</i> pairs	510
Finding the most probable values of <i>r</i> , <i>k</i> , <i>MSY</i> and predicted biomass	511
Bayesian Schaefer analysis	511
Transforming <i>r-k</i> bounds into informative priors	511
Determining a prior for catchability	511
Implementation of the Bayesian Schaefer model	512
<b>Results</b>	<b>512</b>
Results for selecting priors with default rules for <i>r</i> , <i>k</i> , <i>q</i> and biomass	512
Results for simulated stocks with catch and biomass	513
Results for fully assessed stocks	513
Results for simulated stocks with catch and <i>CPUE</i>	514
Results for data-limited stocks	515
<b>Discussion</b>	<b>516</b>
Diversity of examined stocks	516
Priors for <i>r</i> , <i>k</i> and biomass	516
How useful was resilience from FishBase for determining prior ranges of <i>r</i> ?	516
Other options for obtaining prior ranges for <i>r</i>	517
Using maximum catch for obtaining prior ranges for <i>k</i>	517
Is the use of known stock status as prior for biomass circular logic?	517
Data requirements of <i>CMSY</i> compared to other methods	518
Interpretation of the Schaefer equilibrium curve	518
Pragmatic combination of surplus production with recruitment	520
Performance of the Bayesian Schaefer model	521
Understanding the <i>CMSY</i> triangle	521
Performance of <i>CMSY</i>	522
Using <i>CMSY</i> for management of data-limited stocks	522
<b>Conclusions</b>	<b>524</b>
<b>Acknowledgements</b>	<b>524</b>
<b>References</b>	<b>524</b>
<b>Supporting information</b>	<b>526</b>

## Introduction

Most commercially exploited fish stocks in the world lack formal fisheries reference points (Froese *et al.* 2012; Zhou *et al.* 2012), and thus, the degree of exploitation and the status of the stocks are largely unknown. However, legislation in New Zealand (MFNZ 2008), Australia (DAFF 2007), the United States (MSA 2007) and recently also in the European Union (CFP 2013) requires management of all exploited stocks, including those with limited data. Several methods for the assessment of data-limited stocks have been developed (e.g., Cope and Punt 2009; MacCall 2009; Dick and MacCall 2011; Thorson *et al.* 2012; Punt *et al.* 2013; Carruthers *et al.* 2014), and recent reviews of these methods (ICES 2014; Rosenberg *et al.* 2014) have found the

Catch-*MSY* method of Martell and Froese (2013) to be a promising approach. This study revisits the Catch-*MSY* method, addresses its shortcomings, namely the biased estimation of unexploited stock size and productivity, and adds estimation of biomass and exploitation rate. It also addresses a general shortcoming of production models, namely the overestimation of productivity at very low stock sizes (Schnute and Richards 2002; ICES 2014). The predictions of the new method (*CMSY*) are validated against 48 simulated stocks and evaluated against 159 fully or partly assessed real stocks.

## Material and methods

All data files, the R-code of the methods and the figures, a Supplement with detailed presentation of

the methods, default rules for priors and results for 48 simulated and 159 real stocks, and a short text on how to use CMSY are available online for download from <http://oceanrep.geomar.de/33076/>. For convenience, the acronyms and symbols used in this study are summarized in Table 1.

### General description of method


A time series of catches can be viewed as a sequence of yields produced by the available biomass with a given productivity. If two of the three variables, yield, biomass and productivity, are known, then the third can be estimated. Typical production models, such as the one by Schaefer (1954), use time series of catch and abundance to estimate productivity. Instead, the CMSY method presented in this study uses catch and productivity to estimate biomass, providing substantial advancement on the Catch-MSY method of Martell and Froese (2013), which focuses on the estimation of *maximum sustainable yield (MSY)*. CMSY estimates biomass, exploitation rate, MSY and related fisheries reference points from catch data and resilience of the species. Probable ranges for the maximum intrinsic rate of population increase ( $r$ ) and for unexploited population size or carrying capacity ( $k$ ) are filtered with a Monte Carlo approach to detect 'viable'  $r$ - $k$  pairs. A parameter pair is considered 'viable' if the

corresponding biomass trajectories calculated with a production model are compatible with the observed catches in the sense that predicted biomass does not become negative, and is compatible with prior estimates of relative biomass ranges for the beginning and the end of the respective time series. Under these conditions, a plot of viable  $r$ - $k$  pairs typically results in a triangular-shaped cloud in log-space (Fig. 1). The Catch-MSY algorithm (Martell and Froese 2013) was designed to select the most probable  $r$ - $k$  pair as the geometric mean of this distribution. CMSY differs from the Catch-MSY method by searching for the most probable  $r$  not in the centre but rather in the tip region of the triangle. This is based on the underlying principle that defines  $r$  as the maximum rate of increase for the examined population, which should be found among the highest viable  $r$ -values. In other words, a given time series of catches could be explained by a wide range of large stock sizes and low productivity, or by a narrow range of small stock sizes and high productivity, such as in the tip of the triangle (Fig. 1). As  $r$  is defined as maximum net productivity (Schaefer 1954; Ricker 1975), the tip of the triangle is where it should be found.

For verification, the predictions of the CMSY method are compared against simulated data where the 'true' values of parameters and biomass data are known. For evaluation against real-world

**Table 1** Acronyms and symbols used in this study.

Acronym/Symbol	Indicating	Unit
$B$	Biomass, total weight of exploited fish in the water	tonnes
$B_{\text{msy}}$	Biomass capable of producing MSY	tonnes
BSM	Bayesian Schaefer model for estimation of $r$ , $k$ , MSY and $q$	
$C_t$	Catch in a given year	tonnes year <sup>-1</sup>
CMSY	Monte Carlo method for estimating $r$ , $k$ , MSY, biomass and exploitation	
CPUE	Catch per unit effort	$n \text{ h}^{-1}$ or $\text{kg h}^{-1}$
$F_{\text{msy}}$	Rate of fishing mortality compatible with MSY: $F_{\text{msy}} = 0.5 r$	year <sup>-1</sup>
$k$	Parameter of the Schaefer model indicating unexploited stock size	tonnes
$K$	Parameter of the von Bertalanffy somatic growth equation	year <sup>-1</sup>
$M$	Rate of natural mortality	year <sup>-1</sup>
MSY	Maximum sustainable yield: $MSY = r k/4$	tonnes year <sup>-1</sup>
$P_t$	Relative biomass: $P_t = B_t/k$	
$q$	Catchability coefficient: $CPUE_t = q B_t$	
$r$	Maximum intrinsic rate of population increase	year <sup>-1</sup>
$t$	Instant of time; subscript indicating annual values: $C_t$ , $B_t$ or $CPUE_t$	years
$t_{\text{max}}$	Maximum age	years
$t_{\text{gen}}$	Generation time	years
$u$	Exploitation rate: $u_t = C_t/B_t$	
$u_{\text{msy}}$	Exploitation rate that produces MSY at equilibrium	



**Figure 1** Viable  $r$ - $k$  pairs for Pacific Bluefin tuna (*Thunnus orientalis*, Scombridae, BFTuna\_P). The viable  $r$ - $k$  pairs that fulfilled the CMSY conditions are shown in grey. The most probable  $r$ - $k$  pair is marked by the black cross, with indication of approximate 95% confidence limits. The black dots show the estimates of the BSM method, with the white cross indicating the 95% confidence limits. [CMSY\_46eFig 1.R].

fisheries, the predictions of the CMSY method are compared against corresponding parameters and abundance estimates derived from fully or partly assessed stocks, where biomass or *catch-per-unit-effort* (CPUE) data are available in addition to catch data. For this purpose, a Bayesian state-space implementation of the Schaefer model (BSM) is developed, where  $r$ ,  $k$  and  $MSY$  are predicted from catch and abundance data. The basic biomass dynamics are governed by Equation 1:

$$B_{t+1} = B_t + r \left( 1 - \frac{B_t}{k} \right) B_t - C_t \quad (1)$$

where  $B_{t+1}$  is the exploited biomass in the subsequent year  $t+1$ ,  $B_t$  is the current biomass, and  $C_t$  is the catch in year  $t$ .

To account for depensation or reduced recruitment at severely depleted stock sizes, such as predicted by all common stock–recruitment functions (Beverton and Holt 1957; Ricker 1975; Barrowman and Myers 2000), a linear decline of surplus production, which is a function of recruitment, somatic growth and natural mortality (Schnute and Richards 2002), is incorporated if biomass falls below  $\frac{1}{4} k$  (Equation 2).

$$B_{t+1} = B_t + 4 \frac{B_t}{k} r \left( 1 - \frac{B_t}{k} \right) B_t - C_t \mid \frac{B_t}{k} < 0.25 \quad (2)$$

The term  $4 B_t/k$  assumes a linear decline of recruitment below half of the biomass that is capable of producing  $MSY$ .

The BSM was implemented as a Bayesian state-space estimation model (Meyer and Millar 1999; Millar and Meyer 1999), which allowed accounting for variability in both population dynamics (process error) and measurement and sampling (observation error) (Thorson et al. 2014).

The parameters estimated by CMSY and BSM relate to standard fisheries reference points such that  $MSY = r k/4$ , the fishing mortality corresponding to  $MSY$  is  $F_{msy} = 0.5 r$ , the biomass corresponding to  $MSY$  is  $B_{msy} = 0.5 k$  (Ricker 1975; Schaefer 1954) and the biomass below which recruitment may be compromised is half of  $B_{msy}$  (Haddon et al. 2012; Carruthers et al. 2014; Froese et al. 2015).

#### Selection of real stocks and generation of simulated stocks

Altogether 128 fully assessed stocks with biomass data, 28 data-limited stocks with CPUE data and three stocks with less than 9 years of abundance data were used for the evaluation of the CMSY method. Catch and biomass data were extracted from stock assessment documents that were available online or were provided by the respective assessment bodies for the Pacific, North and South Atlantic, the Mediterranean and the Black Sea (see Data S1 for details). In addition to the real stocks, 48 simulated stocks with catch and abundance data were created. The goal was to create a range of biomass scenarios, including strongly as well as

lightly depleted stocks, with monotone stable or monotone changing (i.e. steadily decreasing or increasing) or with alternating biomass trajectories (see Data S1 details).

### CMSY analysis

#### *Determining the boundaries of the r-k space*

To determine prior  $r$ -ranges for the species under assessment, the proxies for resilience of the species as provided in FishBase (Froese et al. 2000; Froese and Pauly 2015) were translated into the  $r$ -ranges shown in Table 2.

Next, a prior range for  $k$  was derived based on three assumptions. First, unexploited stock size  $k$  is larger than the largest catch in the time series, because it is highly unlikely that a fishery finds and catches, in a single year, all individuals of a previously unexploited stock (Vasconcellos and Cochrane 2005; Martell and Froese 2013). Thus, maximum catch in the time series was used to inform the lower bound of  $k$ . Second, the maximum sustainable catch expressed as a fraction of the available biomass ( $F_{msy}$ ) depends on the productivity of the stock. This relationship was accounted for by dividing maximum catch by the upper and lower bound of  $r$  and using these values as the benchmarks for the lower and upper bounds of  $k$ . Third, maximum catch will constitute a larger fraction of  $k$  in substantially depleted rather than lightly depleted stocks. These considerations are summarized in Equation 3 and 4. Suitable ranges for the catch/productivity ratios were determined empirically with simulated data where the true value of  $k$  was known.

$$k_{low} = \frac{\max(C)}{r_{high}}, k_{high} = \frac{4 \max(C)}{r_{low}} \quad (3)$$

where  $k_{low}$  and  $k_{high}$  are the lower and upper bounds of the prior range of  $k$ ,  $\max(C)$  is the

**Table 2** Prior ranges for parameter  $r$ , based on classification of resilience in FishBase (Froese and Pauly 2015).

Resilience	Prior $r$ -range
High	0.6–1.5
Medium	0.2–0.8
Low	0.05–0.5
Very low	0.015–0.1

maximum catch in the time series,  $r_{low}$  is the lower bound of the range of  $r$ -values that the CMSY method will explore and  $r_{high}$  is the upper bound of that range.

$$k_{low} = \frac{2 \max(C)}{r_{high}}, k_{high} = \frac{12 \max(C)}{r_{low}} \quad (4)$$

where variables and parameters are as defined in Equation 3.

Equation 3 was applied to stocks with low prior biomass at the end of the time series and Equation 4 was applied to stocks with high biomass. To reduce the influence of extreme catches, catch data were smoothed by a 3-year moving average.

#### *Setting prior biomass ranges*

To provide prior estimates of relative biomass at the beginning and end of the time series, and optionally also in an intermediate year, one of the possible three broad biomass ranges shown in Table 3 was chosen, depending on the assumed depletion level. This was done automatically by default rules described in the online Supplement. Obvious wrong priors resulting from the default rules, such as setting initial biomass to medium when instead the stock was still lightly exploited or already severely depleted at the beginning of the time series, were noted and subsequently adjusted manually. Thus, the results of this study refer to a scenario where managers are assumed to not have made gross errors in setting broad prior biomass ranges. For example, experts attending the ICES WKLIFE IV and V workshops in Lisbon in October 2014 and 2015 were able to describe stock status and exploitation histories for some of the North Atlantic stocks, which were then translated into the corresponding relative biomass ranges given in Table 2 (ICES 2014, 2015).

#### *Finding viable r-k pairs*

For the detection of viable  $r$ - $k$  pairs, a random  $r$ - $k$  pair is selected from within the prior ranges for  $r$  and  $k$ . Then, a starting biomass is selected from

**Table 3** Default prior biomass ranges relative to  $k$ .

Prior biomass	$B/k$
Low	0.01–0.4
Medium	0.2–0.6
High	0.5–0.9

the prior biomass range for the first year and Equation 1 or 2 is used to calculate the predicted biomass in subsequent years. An  $r$ - $k$  pair is discarded if any of the following conditions applies:

1. The predicted biomass is smaller than 0.01  $k$  (the stock crashes);
2. The predicted biomass falls outside the prior biomass range of the intermediate year;
3. The predicted biomass falls outside the prior biomass range of the final year;

If none of these conditions apply, then the  $r$ - $k$  pair and the trajectory of predicted biomass are considered viable and are stored for analysis. For the purpose of this study, this process was applied to 10 000–200 000 random  $r$ - $k$  pairs, 11–21 start-biomass values and three to six random error patterns for each  $r$ - $k$ -start-biomass combination. To speed up processing, the search for viable  $r$ - $k$  pairs is terminated once more than 1000 pairs are found. For triangles with a thin tip, an additional search is conducted in the tip region.

#### *Finding the most probable values of $r$ , $k$ , MSY and predicted biomass*

CMSY seeks the most probable  $r$ - $k$  pair near the tip of the triangle of viable pairs (Fig. 1). For this purpose, all viable  $r$ -values are assigned to 25–100 bins of equal width in log-space. The 75th percentile of the mid-values of occupied bins is taken as the most probable estimate of  $r$ . This procedure gives equal weight to all occupied bins and reduces the bias caused by the triangular (instead of ellipsoid) shape of the cloud of viable  $r$ - $k$  pairs (compare cloud of probable  $r$ - $k$  pairs estimated by BSM in Fig. 1). Approximate 95% confidence limits of the most probable  $r$  are obtained as 51.25th and 98.75th percentiles of the mid-values of occupied bins, respectively.

The most probable value of  $k$  is determined from a linear regression fitted to  $\log(k)$  as a function of  $\log(r)$ , for  $r$ - $k$  pairs where  $r$  is larger than median of mid-values of occupied bins, with  $\log(4MSY)$  as intercept and with a fixed slope of  $-1$ , based on the rearranged Schaefer model shown in Equation 5. Note that all  $r$ - $k$  pairs on this line have the same intercept and thus give the same value of MSY.

$$MSY = \frac{rk}{4} \rightarrow \log(k) = \log(4MSY) + (-1)\log(r) \quad (5)$$

Approximate 95% confidence limits of  $k$  are obtained by adding the standard deviation of the

residuals of the regression line to the predicted  $k$ -value at the lower confidence limit of  $r$ , and subtracting it from the  $k$ -value predicted for the upper confidence limit of  $r$ . MSY and its 95% confidence limits are obtained as geometric mean of the MSY values calculated for each of the  $r$ - $k$  pairs where  $r$  is larger than the median. Viable biomass trajectories were restricted to those associated with an  $r$ - $k$  pair that fell within the confidence limits of the CMSY estimates of  $r$  and  $k$ . The median of the predicted biomass values for each year was used as the most probable biomass and the 2.5th and 97.5th percentiles were used as indicators of the range that contained 95% of the biomass predictions.

#### **Bayesian Schaefer analysis**

##### *Transforming $r$ - $k$ bounds into informative priors*

For BSM, the uniform  $r$ -ranges shown in Table 2 were translated into prior densities with a central value. An examination of the density of the viable  $r$ -values resulting from CMSY analysis of simulated data was performed using a  $\chi^2$ -test against several standard distributions. The results confirmed that  $r$  is log-normally distributed and suggested that the mean of the  $r$ -ranges in Table 2 provides a reasonable central value. The height of the density function was inversely related to the width of the  $r$ -range, best fit by an *inverse range factor* (*irf*) (Equation 6). The standard deviation of  $r$  in log-space was then described by a uniform distribution between 0.001 *irf* and 0.02 *irf*.

$$irf = \frac{3}{(r_{\text{high}} - r_{\text{low}})} \quad (6)$$

where *irf* is an inverse range factor used in determining the prior density of  $r$  for BSM, and  $r_{\text{high}}$  and  $r_{\text{low}}$  indicate the prior  $r$ -range as defined in Table 2.

The uniform  $k$ -ranges used by CMSY (Equations 3 and 4) were translated into a prior density function by assuming that  $k$  was log-normally distributed and that the mean of the  $k$ -ranges provided a reasonable central value. The standard deviation of the normal distribution in log-space was assumed to be a quarter of the distance between the central value and the lower bound of the  $k$ -range (McAllister et al. 2001).

##### *Determining a prior for catchability*

Data-limited stocks have, by definition, no estimation of biomass but may have, at least for some years, an estimation of stock abundance as CPUE



in units of numbers per hour of fishing or as a biomass index derived from survey catches. Such an abundance index is related to stock biomass by a catchability coefficient  $q$  (Equation 7).

$$CPUE_t = qB_t \quad (7)$$

where  $CPUE_t$  is mean catch per unit effort in year  $t$ ,  $B_t$  is available biomass in year  $t$  and  $q$  is the catchability coefficient. The basic dynamics of the corresponding Schaefer production model for abundance as  $CPUE$  can therefore be expressed in the form of Equation 8.

$$CPUE_{t+1} = CPUE_t + r \left( 1 - \frac{CPUE_t}{qk} \right) CPUE_t - qC_t \quad (8)$$

where variables and parameters are as defined in Equations 1 and 7. Priors for  $q$  were derived from the Schaefer equilibrium equation for catch (Equation 9).

$$Y = rB \left( 1 - \frac{B}{k} \right) \quad (9)$$

where  $Y$  is the equilibrium yield for any given biomass  $B$ , and other parameters are as defined in Equation 1.

Setting  $B/k = 0.5$ ,  $B = CPUE/q$  and  $Y = \text{catch}$  gives  $q = 0.5 r CPUE/C$  for  $MSY$ -level catch and biomass. Setting  $B/k = 0.25$  for half  $MSY$ -level biomass gives  $q = 0.75 r CPUE/C$ . Suitable multipliers for low and high biomass and prior  $r$ -ranges were derived empirically from simulated data. For stocks with high recent prior biomass, priors for  $q$  were derived as shown in Equation 10 and 11.

$$q_{\text{low}} = \frac{0.25 r_{\text{pgm}} CPUE_{\text{mean}}}{C_{\text{mean}}} \quad (10)$$

where  $q_{\text{low}}$  is the lower prior for the catchability coefficient for stocks with high recent biomass,  $r_{\text{pgm}}$  is the geometric mean of the prior range for  $r$ ,  $CPUE_{\text{mean}}$  is the mean of catch per unit effort over the last 5 or 10 years, and  $C_{\text{mean}}$  is the mean catch over the same period.

$$q_{\text{high}} = \frac{0.5 r_{\text{high}} CPUE_{\text{mean}}}{C_{\text{mean}}} \quad (11)$$

where  $q_{\text{high}}$  is the upper prior for the catchability coefficient for stocks with high recent biomass,  $r_{\text{high}}$  is the upper prior range for  $r$  and all other variables are as defined in Equation 10.

For stocks with low recent prior biomass, the multipliers were changed from 0.25 to 0.5 for  $q_{\text{low}}$  and from 0.5 to 1.0 for  $q_{\text{high}}$ . Mean catch and  $CPUE$

were taken over the last 5 years for species with medium and high resilience or over the last 10 years for species with low or very low resilience. For the Bayesian implementation of the Schaefer model, the  $q$ -range was translated into a prior density function by assuming that  $q$  was log-normally distributed and that the mean of the log  $q$ -range provided a reasonable central value, with a standard deviation assumed to be a quarter of the distance between the central value and  $q_{\text{low}}$  (McAllister *et al.* 2001). The implementation of BSM for  $CPUE$  data used the same settings as applied to observed or simulated biomass (see below). If less than 9 years of  $CPUE$  data were available, the Schaefer model was not fit. Instead,  $CPUE$  was plotted on a second  $y$ -axis in the plot of biomass predicted by CMSY (see example in Fig. 6).

#### Implementation of the Bayesian Schaefer model

The state-space model implementation of the BSM (Millar and Meyer 1999) for catch and biomass and for catch and  $CPUE$  are included in the CMSY R-code, which is available as part of the online Supplement. The JAGS software (Plummer 2003) was used for sampling the probability distributions of the parameters with the Markov chain Monte Carlo method. To facilitate mixing of the Gibbs samples, annual biomass was expressed relative to the unexploited biomass with  $P_t = B_t/k$  (Meyer and Millar 1999). Basic parameter settings included three sampling chains with a chain length of 60 000 steps each and with a burn-in phase of 30 000 steps. For the analysis of output, only every 10th value was used to reduce autocorrelation. All posterior parameter estimates were assumed to be approximately log-normally distributed, with the median used as the central value and 95% confidence intervals approximated by the 2.5th and 97.5th percentiles to find values at which test statistics attain less than 0.05 significance (Gelman *et al.* 1995; McAllister *et al.* 2001; Owen 2013).

## Results

### Results for selecting priors with default rules for $r$ , $k$ , $q$ and biomass

To determine the prior ranges for  $r$ , resilience categories (very low, low, medium, high) at the species level were used from [www.fishbase.org](http://www.fishbase.org) for fishes and were selected manually for the four stocks of invertebrates. These categories, combined with

catch data and prior biomass ranges, led to the detection of viable  $r$ - $k$  pairs in 140 of 159 stocks (88%). In 19 stocks, the resilience category was changed manually to an adjacent lower or upper category in order to find viable  $r$ - $k$  pairs. In about half of these cases (nine of 19), the change was to an upper resilience category, that is there was no apparent bias in the default rules towards lower or higher categories.

The rules for deriving the prior ranges for  $k$  from maximum catch and prior  $r$  were sufficient in all cases; so that no manual adjustments were done. Priors for catchability  $q$  were derived from equilibrium reasoning and recent catches (Equations 9–11). The prior ranges included the ‘true’ value of  $q$  in 19 of 24 simulated stocks (79%, see Table 8).

To determine prior biomass ranges for the start and end of the time series, and for an optional intermediate year, default rules were used as described in the online Supplement. The resulting prior biomass ranges were compatible with observed biomass or abundance in 92 of 159 stocks (58%). Initial biomass was manually corrected in 14 stocks (9%), intermediate biomass in 11 stocks (9%) and final biomass in 54 stocks (34%). See data on resilience and biomass priors in AllStocks\_ID20.xlsx in the online Supplement material.

#### Results for simulated stocks with catch and biomass

The CMSY and BSM methods were applied to simulated catch and biomass data where the ‘true’ parameter values were known. Tabulated results and detailed analyses for every stock are available

in the online Supplement (Tables S3, S4 and Appendix I in Data S1). In most simulated stocks (75–96%, depending on the parameter, see Table 4), the 95% confidence limits of the estimates by CMSY and BSM included the true value used in the simulations and were thus not significantly different from the ‘true’ values (Smith 1995). Results for CMSY and BSM were very similar. Of the six scenarios where CMSY estimates of  $r$  did not include the ‘true’ value, four had high final biomass. Similarly, all four scenarios where BSM estimates did not include the ‘true’ value had high final biomass.

A comparison of CMSY and BSM estimates vs. ‘true’ values for  $MSY$ ,  $r$ ,  $k$ , last biomass and last exploitation rate showed that the median ratios and the ranges that contain 90% of the estimates were very similar for both methods (Table 5). The median ratios were close to unity, with maximum deviations of 0.80 and 1.06. The 5th–95th percentile ranges, which contain 90% of the estimates, included the expected ratio of 1.0 in all cases and were bracketed by the ratios 0.69 and 1.62 for BSM and 0.20 and 5.71 for CMSY. The latter strong deviation referred to relative biomass in the simulated stock HL\_M. In this case, the ‘true’ value in the final year was 0.002  $k$ , whereas the CMSY estimate was 0.109  $k$ . The deviation is caused by the default prior for low biomass of 0.01–0.4  $k$ , which excludes the ‘true’ biomass.

#### Results for fully assessed stocks

The CMSY and BSM methods were applied to 128 real stocks for which catch and biomass data were available from recent stock assessments. Detailed

**Table 4** Number and percentage of 24 simulated stocks which contain the true parameter value within the 95% confidence limits of the respective CMSY or BSM estimate. The acronyms of the stocks where the confidence limits do not include the true value are presented in the column ‘Missed stocks’ (Appendix I in Data S1). [SimCatchResults\_6.xlsx].

Parameter	True value included in 95% confidence limits	Per cent	Missed stocks
$r$ CMSY	18	75.0	HH_M, HLH_M, LH_H, LH_VL, LHL_M, LHL_VL
$r$ BSM	19	79.2	HH_M, HL_H, HLH_M, LH_H, LH_M
$k$ CMSY	20	83.3	HLH_M, LH_H, LH_VL, LHL_M
$k$ BSM	20	83.3	HH_M, HL_M, LH_H, LHL_VL
$MSY$ CMSY	23	95.8	LL_M
$MSY$ BSM	18	75.0	HH_M, HL_H, HL_M, HLH_VL, LH_M, LHL_VL
Last $B/k$ CMSY	22	91.7	HL_M, LHL_M



**Table 5** Comparison of the estimation of reference points with the BSM and with the CMSY method, applied to 24 simulated data sets where the true values of the reference points were known. [SimCatchResults\_6.xlsx].

Ratio	Median	5th percentile	95th percentile
$MSY_{BSM}/true\ MSY$	1.01	0.69	1.60
$MSY_{CMSY}/true\ MSY$	1.02	0.53	1.62
$r_{BSM}/true\ r$	1.05	0.93	1.31
$r_{CMSY}/true\ r$	1.06	0.59	1.73
$k_{BSM}/true\ k$	0.98	0.79	1.29
$k_{CMSY}/true\ k$	1.00	0.58	1.80
Last $B/k_{CMSY}/last\ true\ b/k$	1.10	0.80	5.71
Last $u_{CMSY}/last\ true\ u$	0.85	0.20	1.72
Last $u/u_{msy}\ CMSY/last\ true\ u/u_{msy}$	0.80	0.22	1.57

analyses are available for every stock in Appendix II and summarized in Tables S5 and S6 of Data S1. For most stocks (71–79%, depending on the parameter, see Table 6), the 95% confidence limits of the CMSY estimates included the most probable BSM estimate, indicating good agreement between the methods (Smith 1995). In 5–16% of the stocks, the confidence limits of both methods did not overlap, indicating that the predictions were significantly different (Knezevic 2008).

A comparison of CMSY and BSM estimates for  $MSY$ ,  $r$ ,  $k$ , final biomass and exploitation rate in the final year shows that the median ratios were generally close to 1.0, with maximum deviations of 0.92 and 1.08. The 5th–95th percentile ranges always included unity and were bracketed by the

**Table 6** Numbers and percentages of 128 real stocks, where the 95% confidence limits of the CMSY estimate include the most probable estimate of BSM, indicating good agreement, and where the confidence limits of both methods did not overlap, indicating results that are significantly different. [AllStocks\_Results\_6.xlsx].

Parameter	BSM estimate included n/%	No overlap n/%
$r_{CMSY}$	101/78.9	14/10.9
$k_{CMSY}$	101/78.9	20/15.6
$MSY_{CMSY}$	97/75.8	6/4.7
Last relative $B_{CMSY}$	91/71.1	13/10.2

ratios 0.47 and 2.16 for  $r$ ,  $k$  and  $MSY$ , but were wider (0.34–2.91) for the last year's CMSY estimates of biomass and exploitation rate compared to observed data (Table 7).

### Results for simulated stocks with catch and CPUE

The CMSY and BSM methods were applied to simulated catch and  $CPUE$  data where the true parameter values were known. Tabulated results and detailed analyses for every stock (Appendix III in Data S1) are available in the online Supplement (Tables S7, S8 and Appendix III in Data S1). The use of  $CPUE$  rather than biomass did not affect the CMSY results, because neither biomass nor  $CPUE$  are used by CMSY. The Bayesian implementation of the Schaefer model for catch and  $CPUE$  data required the additional estimation of catchability  $q$  to transform  $CPUE$  into biomass. The 95% confidence limits of  $q$  estimated by BSM included the true value in 33% of the cases (Table 8), however, with mostly narrow and only three substantial misses (Table S8 in Data S1). For the other parameters ( $r$ ,  $k$ ,  $MSY$ , last biomass), the 95% confidence limits estimated by CMSY included the true values in 67–96% and for BSM in 50–88% of the cases (Table 8). The lower success rate of BSM is due to its confidence limits being generally narrower than for CMSY.

A comparison of CMSY estimates vs. true values for  $MSY$ ,  $r$ ,  $k$  and last biomass showed median ratios close to 1.0 (0.99–1.15) with ranges that contained 90% of the estimates from 0.51 to 5.05.

**Table 7** Comparison of the estimation of reference points with the CMSY method and with BSM, applied to 128 fully assessed stocks. [AllStocks\_Results\_6.xlsx].

Ratio	Median	5th percentile	95th percentile
$r_{CMSY}/r_{BSM}$	0.97	0.58	1.62
$k_{CMSY}/k_{BSM}$	1.00	0.47	2.16
$MSY_{CMSY}/MSY_{BSM}$	1.02	0.65	1.58
Last $B_{CMSY}/last\ observed\ B$	0.93	0.34	2.68
Last $B/k_{CMSY}/last\ observed\ B/k$	0.92	0.47	1.53
Last $u_{CMSY}/last\ observed\ u$	1.08	0.37	2.91
Last $u/u_{msy}\ CMSY/last\ observed\ u/u_{msy}$	1.06	0.48	2.45

**Table 8** Numbers and percentages of 24 stocks with simulated catch and CPUE data, where the 95% confidence limits of BSM or CMSY include the 'true' value. The acronyms of the stocks where the confidence limits of the respective parameters do not include the 'true' value are given in the column 'Missed stocks' (Appendix III in Data S1). [SimCatchCPUE\_Results\_6.xlsx].

Parameter	True value included in 95% confidence limits	Per cent	Missed stocks
Prior for $q$	19	79.2	HH_M, HLH_M, HLH_VL, LH_H, LH_VL
$q$ BSM	8	33.3	many
$r$ CMSY	18	75.0	HH_M, HLH_M, LH_H, LH_VL, LHL_M, LHL_VL
$r$ BSM	12	50.0	many
$k$ CMSY	20	83.3	HLH_M, LH_H, LH_VL, LHL_M
$k$ BSM	13	54.2	many
$MSY$ CMSY	23	95.8	LL_M
$MSY$ BSM	21	87.5	HLH_VL, LH_VL, LL_L
Last $B$ CMSY	16	66.7	many
Last $B/k$ CMSY	22	91.7	HL_M, LHL_M

The last number refers to relative biomass in the simulated stock HL\_M, where the 'true' value in the final year was 0.002  $k$ , whereas the CMSY estimate was 0.096  $k$ . The deviation is caused by the default prior for low biomass of 0.01–0.4  $k$ , which excludes the 'true' biomass. The BSM method for data-limited stocks had median ratios of 0.93–1.07 and 90% ranges from 0.39 to 6.22, where the latter value stems from the three estimates of  $q$  which differed substantially from the 'true' value (Table 9).

### Results for data-limited stocks

Altogether 31 data-limited stocks were analysed with the CMSY method. Twenty-eight of these stocks had sufficiently long ( $\geq 9$  years) time series of abundance available so that BSM could also be applied and CMSY and BSM estimates could be compared. Detailed analyses for every stock as well as summary tables are available in the online Supplement (Appendix IV and Tables S9, S10 in Data S1). In most stocks, the 95% confidence limits of the CMSY estimates for  $r$ ,  $k$  and  $MSY$  included the

**Table 9** Comparison of the estimation of reference points with BSM and with CMSY, applied to 24 data sets with simulated CPUE, where the 'true' values of the reference points are known. [SimCatchCPUE\_Results\_6.xlsx].

Ratio	Median	5th percentile	95th percentile
$q$ BSM/true $q$	1.07	0.61	6.22
$MSY$ BSM/true $MSY$	0.97	0.39	2.03
$MSY$ CMSY/true $MSY$	1.02	0.51	1.61
$r$ BSM/true $r$	1.02	0.76	1.56
$r$ CMSY/true $r$	1.03	0.63	1.74
$k$ BSM/true $k$	0.93	0.36	1.86
$k$ CMSY/true $k$	0.99	0.60	1.64
Last $B$ CMSY/last true $B$	1.15	0.60	5.05

**Table 10** Numbers and percentages of the 28 data-limited stocks where the 95% confidence limits of the CMSY estimates include the most probable estimate of BSM, and where the confidence limits of both methods did not overlap, indicating results that are significantly different. [CPUEStocks\_Results\_6.xlsx].

Parameter	BSM estimate included $n/\%$	No overlap $n/\%$
$r$ CMSY	27/96.4	1/3.6
$k$ CMSY	27/96.4	27/3.6
$MSY$ CMSY	25/89.3	27/3.6
Last $B$ CMSY	20/71.4	24/14.3
Last relative $B$ CMSY	19/67.9	24/14.3
$u$ CMSY	20/71.4	24/14.3

most probable BSM estimate. Depending on the parameter, the 95% confidence limits of CMSY included the BSM estimate in 68–96% of the stocks (Table 10), suggesting good agreement between the methods (Smith 1995). The confidence limits of both methods did not overlap in 4–14% of the stocks, indicating that the respective estimates were significantly different (Knezevic 2008). A comparison of CMSY and BSM estimates for  $MSY$ ,  $r$ ,  $k$ , final biomass and exploitation rate in the final year showed that the median ratios and the ranges containing 90% of the estimates were similar for both methods (Table 11). The median ratios were close to 1.0, with maximum deviations of 1.05 and 1.37. The 5th–95th percentile ranges were bracketed by the ratios 0.63 and 1.68 for  $r$ ,  $k$  and  $MSY$ , but were wider (0.51–

**Table 11** Comparison of the estimated reference points from the CMSY method and from the Bayesian Schaefer model, applied to 28 data-limited stocks. [CPUEStocks\_Results\_6.xlsx].

Ratio	Median	5th percentile	95th percentile
$r$ CMSY/ $r$ BSM	1.06	0.75	1.17
$k$ CMSY/ $k$ BSM	1.07	0.63	1.60
MSY CMSY/MSY BSM	1.05	0.70	1.68
Last $B$ CMSY/last $B$ BSM	1.37	0.62	7.84
Last $B/k$ CMSY/last observed $B/k$	1.26	0.51	12.8

12.8) for the final-year estimates of biomass. The strong deviation was caused by two stocks (cod-rock, smn-sp) where biomass was severely depleted to less than 1% of unexploited stock size, while the default biomass prior used by CMSY still assumed 1–40% of unexploited biomass.

## Discussion

### Diversity of examined stocks

We collected 159 time series of catch and abundance from various sources, including stocks from the North and South Pacific, North and South Atlantic, the Caribbean, the Mediterranean and the Black Sea. Four stocks belonged to two species of crustaceans and the remaining stocks belonged to species of marine fishes, including five elasmobranchs. Over two-third of the species were demersal. Other fishes consisted of small pelagic and highly migratory species such as tunas and billfishes. Species were distributed mostly in temperate climate zones, several in subtropical zones, some in polar regions, and only one in the tropics. Deep-sea fishes were represented by eleven species. Very low- to medium-resilience categories were represented by 14 or more stocks each, but only four stocks fell into the high-resilience category. Several species were represented by more than one and up to 12 different stocks. Several time series used in the analysis started as early as 1930 and most reached until 2012. The highest catch was reported for blue whiting (*Micromesistius poutasou*, Gadidae) with 2.4 million tonnes in the Northeast Atlantic in 2004, and the largest stock

size was reported for Eastern Bering Sea pollock (*Theragra chalcogramma*, Gadidae) with an estimated biomass of 13.1 million tonnes in 1995. This selection of stocks was reasonably complete and representative for the Northeast Pacific and the North Atlantic, but by no means complete or representative for the other areas or for global fisheries. However, the selection of stocks covered a wide range of the diversity of commercial species, suggesting that the methods and default rules for priors used in this study are broadly applicable.

### Priors for $r$ , $k$ and biomass

*How useful was resilience from FishBase for determining prior ranges of  $r$ ?*

The resilience categories from FishBase (Froese et al. 2000; Froese and Pauly 2015) gave prior ranges of  $r$  that led to similar and therefore presumably reasonable CMSY and BSM fits for 88% of the stocks. For the remaining stocks, reasonable fits were only obtained if the next higher or lower resilience category was chosen. The even distribution of these corrections between lower and upper resilience suggests that these species were intermediate to the available resilience categories. Also, for some species, different stocks required different categories. For example, the two examined stocks of surmullet (*Mullus surmulletus*, Mullidae) gave reasonable fits for CMSY and BSM only when resilience was set medium for the North Sea stock (mur-347d) and high for the Central Mediterranean stock (mullsur\_gsa1516). Similarly, of the eight examined stocks of haddock (*Melanogrammus aeglefinus*, Gadidae), seven gave reasonable fits with the FishBase category of medium resilience, whereas the stock from Georges Bank (Haddock\_GB) only gave a reasonable fit when resilience was set to low. More generally, if the prior  $r$ -range is set too high, CMSY is unlikely to find viable  $r$ - $k$  pairs; if the prior  $r$ -range is set too low, the  $r$ - $k$  space is likely to be flooded with viable  $r$ - $k$  pairs pressing against the upper bound of  $r$ . Thus, while the resilience categories from FishBase provided a good starting point for prior ranges of  $r$ , users of the CMSY and BSM methods should carefully consider all available information and then select the most suitable prior range of  $r$  for the stock in question, independent of the fixed ranges used for the purpose of this study (Table 2).

*Other options for obtaining prior ranges for r*

In the context of the Schaefer model, half of the maximum intrinsic rate of population increase  $r$  equals the rate of fishing mortality  $F_{\text{msy}}$  that is compatible with MSY (Ricker 1975).  $F_{\text{msy}}$  itself is closely related to the rate of natural mortality  $M$  (Zhou et al. 2012; Froese et al. 2016a,b). Thus, if estimates of  $F_{\text{msy}}$  or  $M$  are available for other stocks of the respective species, or such estimates are available for similar species in the same area, then these estimates can be used to determine prior ranges for parameter  $r$ . Jensen (1996) suggested an evolutionary relationship between natural mortality and the somatic growth parameter  $K$ , such that  $K = 2/3 M$ . Kenchington (2014) examined 29 estimators of natural mortality and concluded that  $M = 1.5 K$  or Pauly's (1980) empirical estimator based on growth parameters and temperature or maximum age ( $t_{\text{max}}$ ) with  $M = 4.3/t_{\text{max}}$  can provide useful estimates. Generation time ( $t_{\text{gen}}$ ) is also a strong predictor for  $r$  (Myers et al. 1999; Froese et al. 2000; McAllister et al. 2001). FishBase (Froese and Pauly 2015) has compiled mortality estimates for several hundred species and maximum age and growth studies for several thousand species, including most commercial species. These published parameter estimates can be used to establish prior ranges for  $r$ . Equation 12 summarizes the approximate relation between  $r$  and other life history parameters that may be more easily available.

$$r \approx 2F_{\text{msy}} \approx 2M \approx 3K \approx 3/t_{\text{gen}} \approx 9/t_{\text{max}} \quad (12)$$

*Using maximum catch for obtaining prior ranges for k*

There is no simple predictor for the unexploited size of a population because estimates of abundance typically start well after fishing has substantially reduced stock size. However, it is highly unlikely that a fishery catches the whole stock in a single year, and thus, it is safe to assume that the maximum catch in a time series will be smaller than the unexploited stock size  $k$ . Therefore, maximum catch modulated by productivity was used as a reference point for determining prior ranges of  $k$ . It can be argued that catch data are the main input of CMSY and that catch must therefore be treated as unknown when establishing priors (e.g. Kruschke 2011). However, the prior knowledge used here is not the catch itself but the knowledge that the unknown  $k$  must be larger than the maximum catch and that

populations with high productivity can sustain larger maximum catches relative to  $k$ . For the wide range of species and stocks examined in this study, the prior range for  $k$  was suitable for CMSY and BSM analyses and thus seems fit for general use.


*Is the use of known stock status as prior for biomass circular logic?*

Broad estimates of relative biomass at the beginning and the end of the time series of catches are required inputs for CMSY. For the purpose of this study, these prior biomass ranges were set by default rules as low, medium or high (Table 2). The default rules gave satisfactory results in about 2/3 of the stocks. But in 34% of the stocks, the prior for final biomass had to be corrected to include, or nearly include, observed abundance. Thus, one-third of the analyses presented in this study refer to a case where experts had selected prior biomass ranges that were compatible with the true status of the stock. While this may sound like circular logic, independent knowledge about stock status often exists and its inclusion in the analysis is then mandatory in a Bayesian context (Gelman et al. 1995; Kruschke 2011).

For example, it is well-known that the North Sea herring stock (*Clupea harengus*, Clupeidae, her-47d3) was in reasonably good state in the 1950s, collapsed in the 1970s and has recovered in recent years. FishBase gives the resilience of this species as medium. This very general information combined with the time series of catches suffices to produce CMSY estimates of  $r$ ,  $k$ , MSY and trajectories of biomass and exploitation that are similar to the respective estimates produced by BSM and by regular stock assessment (Fig. 2).

Other examples of well-known stock status histories are Georges Bank cod (*Gadus morhua*, Gadidae, Cod\_GB), which was overfished in the 1980s, collapsed in the 1990s and has not recovered since, or Arctic cod (*Gadus morhua*, Gadidae, cod-arct), which was abundant in the early 1950s, was near collapse in 1990 and recovered in recent years.

Some preliminary testing of the sensitivity of CMSY to incorrectly set prior biomass ranges was carried out at the WKLIFE V workshop (ICES 2015). As expected, especially long time series were able to recover from incorrectly set initial or intermediate prior biomass ranges, but not from incorrectly set final biomass ranges, because, by



**Figure 2** Results of CMSY analysis for North Sea herring (her-47d3). Panel (a) shows the biomass trajectory predicted by CMSY in bold, together with the 2.5th and 97.5th percentiles. The dashed curve is the observed biomass, scaled by  $k$  as estimated by BSM. The dotted vertical lines indicate the prior ranges for relative biomass. The dashed horizontal line indicates  $B_{msy} = 0.5 k$  and the dotted horizontal line indicates half  $B_{msy}$  as the border below which recruitment may be reduced. Panel (b) shows the relative exploitation rate  $u/u_{msy}$  as estimated by CMSY in bold. The dashed curve shows the observed ratio of catches over total biomass, scaled by  $0.5 r$  as estimated by BSM. [CMSY\_46eFig 2\_2.R].

design, final biomass estimates outside the prior range are discarded by the CMSY algorithm.

However, similar constraints also apply to many data-limited and data-moderate assessments based on conventional age-structured models (Ludwig and Walters 1989; Mangel *et al.* 2013). Firstly, in the absence of complete catch time series that date back to the onset of the fishery, it is typically necessary to enforce reasonable initial or current biomass estimates through informative prior or penalties (e.g. Punt and Hilborn 1997). Secondly, key parameters such as the natural mortality or the steepness of the assumed spawner-recruitment relationship are often fixed or heavily constrained, due to limited information in the available data (Lee *et al.* 2011, 2012). This means that other important reference points and thus the outcome of the assessment may be determined *a priori* (Mangel *et al.* 2013). By comparison, CMSY admits broad uncertainty in resilience and productivity and may therefore be more robust to model misspecifications (Thorson *et al.* 2014). In summary, the inclusion of independent knowledge about stock status is neither unique to CMSY nor circular logic but mandatory in a Bayesian-type analysis. CMSY is most sensitive to incorrect setting of final prior biomass, but experts and stakeholders for a given stock are likely to have reasonably correct knowledge about current stock status.

#### Data requirements of CMSY compared to other methods

In comparison with other methods proposed for data-limited stock assessment, the requirements of

CMSY (catch, qualitative resilience and qualitative stock status) appear modest. For example, the DCAC method (MacCall 2009), which estimates a sustainable catch-level below  $MSY$ , requires catch, relative depletion,  $M$  and  $F_{msy}/M$  as inputs. The DB-SRA method (Dick and MacCall 2011), which estimates biomass,  $MSY$ ,  $B_{msy}$  and exploitation rate, requires catch, relative depletion,  $M$ ,  $F_{msy}/M$ ,  $B_{msy}/B_0$  and age at maturity as inputs. The COM-SIR method (Vasconcellos and Cochrane 2005), which estimates stock status, production and exploitation rates, requires catch, priors for  $r$  and  $k$ , relative bioeconomic equilibrium and increase in harvest rate over time as inputs. The SSCOM method (Thorson *et al.* 2013), which predicts stock status and productivity, requires catch, priors for unexploited biomass, initial effort and parameters of an effort-dynamics model. Reviews of these and other data-limited methods concluded that the Catch- $MSY$  method (Martell and Froese 2013), of which CMSY is an advanced implementation, performed best with respect to proportional error in predictions (ICES 2014; Rosenberg *et al.* 2014).

#### Interpretation of the Schaefer equilibrium curve

CMSY and BSM as implemented here do not fit a parabola to yield and biomass data, which requires the assumption of equilibrium conditions, but rather search for the  $r$ - $k$  pair that best fits the time series of available data and prior information. Plotting catch and biomass data against an




equilibrium surplus production curve often looks unconvincing and may lead to erroneous estimates of  $MSY$  and  $B_{msy}$  if the stock was not in a state of equilibrium (Punt 2003). This is depicted in Fig. 3a, which shows CMSY and BSM estimates as well as ‘true’ data points for a simulated stock of low resilience with increasing biomass. All points fall below the equilibrium curve, because catches were consistently less than surplus production throughout the time series, which allowed the stock to increase. Also, the non-overlap of the CMSY estimates with the ‘true’ points and the BSM estimates suggests that the CMSY fit for this stock is not particularly good. By contrast, Fig. 3b shows a simulated stock with high resilience and a declining biomass pattern, including phases of increase, decrease and equilibrium, as indicated by the position of points below, above and near the equilibrium curve, respectively. The overlap of points suggests good agreement between CMSY and BSM estimates and ‘true’ parameter values.

Production models with equilibrium curves of different shapes have been proposed, but they all share the same anchor points, because zero biomass produces zero yield at the one end, and zero yield results in unexploited biomass at the other end. All models have an intermediate maximum at similar absolute catch and biomass values, as forced by the data (Sparre and Venema 1998; Froese *et al.* 2011; Thorson *et al.* 2012). Only the estimation of unexploited biomass and therefore the relative position of the biomass that can

produce  $MSY$  changes with the choice of the production model, from  $0.37 B/k$  with the Fox (1970) model to  $0.5 B/k$  with the Schaefer (1954) model, and somewhere in between with the Pella and Tomlinson (1969) model, depending on the shape parameter of that model. The parabolas shown in Fig. 3 are from a Schaefer model, representing surplus production derived from the first derivative of the logistic model of population growth. The Schaefer model has fewer assumptions and is more conservative than the other models (it predicts lower equilibrium catch at low biomass, see Figure A1 in the Supporting Information of Froese *et al.* 2011) and was therefore chosen for the implementation of CMSY in this study. The indentation of the parabolas shown in Fig. 3 at stock sizes below  $0.25 k$  results from the inclusion of a stock–recruitment model which assumes reduced recruitment at low stock sizes (see next section).

Equilibrium conditions may arise when the same catch is taken over an extended period of time of at least one generation (Kawasaki 1980; Caddy 1984). Thus, persistent deviations in the annual points from the equilibrium curve are more likely to be found in species with low resilience and long generation time, such as in the simulated species with low resilience in Fig. 3a. In contrast, a wide and more evenly distributed scattering of points can be expected in species with high resilience and short generation times, such as simulated in Fig. 3b. In summary, while the




**Figure 3** CMSY output showing catch relative to  $MSY$  over biomass relative to unexploited stock size for two simulated stocks. The open circles indicate true data, the black dots indicate estimates by BSM, and the grey dots indicate estimates by CMSY. Panel (a) shows results for a simulated stock with increasing biomass and low resilience (see LH\_L in Appendix I of Data S1). Panel (b) shows results for a simulated stock with declining biomass for a species with high resilience (see HL\_H in Appendix I of Data S1). The indentation of the parabolas below  $0.25 k$  (half of  $B_{msy}$ ) results from the inclusion of a stock–recruitment model which assumes reduced recruitment at low stock sizes. [CMSY\_46eFig 3\_2.R].

equilibrium curve is not suitable for parameter estimation in most situations, it is still useful for understanding the status of the stock and for comparing CMSY and BSM estimates.

#### Pragmatic combination of surplus production with recruitment

Production models have been criticized for not taking into account the widely observed reduction in recruitment at low population sizes. Instead, these models assume an increase in the biomass growth rate  $dB/dt$  as biomass approaches zero (Schnute and Richards 2002). In earlier versions of CMSY, this increase in productivity with decreases in biomass led to an overestimation of final biomass in depleted stocks (see respective warnings in ICES 2015, Annex 3). Schnute and Richards (2002) propose to solve the general problem by combining the production model with a recruitment function, but their solution consists of eight interconnected equations with more than eight additional parameters to be estimated. Here, we choose a much simpler approach, assuming a generic stock–

recruitment function with constant recruitment above  $0.25 k$  and linear decline of recruitment below that threshold, towards zero recruitment at zero biomass. Such a hockey-stick model of recruitment has been proposed by Barrowman and Myers (2000), and a threshold around half of  $B_{msy}$  has been widely adopted as a limit reference point for recruitment overfishing (Beddington and Cooke 1983; Myers *et al.* 1994; Punt *et al.* 2013; Caruthers *et al.* 2014; Froese *et al.* 2016a,b). A hockey-stick function is combined here with the production model by introducing a multiplier which decreases linearly from 1 to zero at biomass below  $0.25 k$ . This multiplier is assumed to reduce the unknown component that recruitment provides to surplus production (Equation 2). This new ‘surplus production and recruitment’ model is used in CMSY and BSM and gives more realistic estimates of  $r$  and  $k$  in stocks with extended periods of severely depleted biomass. It also removes the bias in CMSY estimates of final biomass in severely depleted stocks (see YTFlo\_MA and her-3a22 in Fig. 4, with reasonable predictions of final biomass despite severe depletion). Note that the reduction



**Figure 4** CMSY predictions of relative biomass  $B/k$  (bold curve) with 2.5th and 95th percentiles (thin curves) compared to observed biomass (dashed curve) scaled by the respective BSM estimate for  $k$  for (a) North Atlantic swordfish (Swordfish\_NA), (b) Arctic cod (cod-arct), (c) Northwest Atlantic yellow tail flounder (*Limanda ferruginea*, Pleuronectidae, YTFlo\_MA) and (d) Western Baltic herring (*Clupea harengus*, Clupeidae, her-3a22). The horizontal lines emphasize  $B_{msy} = 0.5 k$  and  $0.5 B_{msy} = 0.25 k$ . The dotted vertical lines indicate the prior estimates of biomass. [CMSY\_46eFig 4\_2.R].

in recruitment at very low stock sizes ( $B/k < 0.25$ ) also means that  $F_{\text{msy}} = \frac{1}{2} r$  is not applicable anymore and instead  $F_{\text{msy}} = \frac{1}{2} r \frac{4}{B/k}$  should be used for management advice.

### Performance of the Bayesian Schaefer model

Dynamic production models, such as the implementation of the Schaefer model used in this study, require time series of catch and abundance as inputs and thus do not count as data-limited methods. However, as CMSY is a simplified Bayesian implementation of a data-limited production model, it seems appropriate to compare CMSY results with the results of a full Bayesian implementation of a surplus production estimation model, rather than with results obtained from various stock assessment methods with different assumptions and often unavailable levels of uncertainty. The Bayesian Schaefer model applied in this study (BSM) is similar to previous implementations (e.g. Meyer and Millar 1999; McAllister *et al.* 2001; Vasconcellos and Cochrane 2005; Thorson *et al.* 2013), but differs in its emphasis on informative priors for  $k$ , based on maximum catch modulated by productivity, for  $q$ , based on equilibrium catch, for  $r$ , based on more complex modelling of the distribution of  $r$ , and for relative biomass ranges, based on default rules or expert opinion. A state-space model implementation was chosen because explicit modelling of process error and observation error has been shown to result in more realistic posterior distributions of parameters (Ono *et al.* 2012). The resulting predictions for  $r$ ,  $k$  and  $MSY$  were close to the 'true' values of the simulated data sets and  $r/2$  was reasonably close to working group estimates of  $F_{\text{msy}}$  in 82% of the stocks with available data (Data S1).

In summary, BSM performed well when compared with 'true' values of simulated stocks. It could be fitted to all real stocks with at least 9 years of abundance data and produced parameter estimates that were comparable with available working group estimates. Thus, the BSM parameter estimates were chosen as benchmarks for the evaluation of CMSY when applied to real stocks where true parameter values are unknown. Of course, like any production model, BSM will provide unrealistic results if one or more of its key assumptions are violated, caused, for example, by environmental regime shifts, dramatic changes in

the productivity or size-structure of the stock, or major changes in catchability. Also, in stocks that are lightly exploited such as the simulations ending in high biomass, the interplay between catch and biomass contains less information about productivity and estimates of  $r$  will be less reliable (Table 4).

### Understanding the CMSY triangle

The typical triangular shape of viable  $r$ - $k$  points in log-space (Fig. 1) has a biological basis. The given time series of catches may have been produced either by a large population with low to medium productivity or by a small population with high productivity. This relation is reflected by the well-defined lower bound of the  $r$ - $k$  triangle, which represents for a given  $r$  the lowest  $k$  that is compatible with the catches and the biomass priors. The slope of this lower border is typically a bit flatter than the slope ( $-1$ ) of the line of  $r$ - $k$  pairs resulting in the same value of  $MSY$  (Equation 5). The upper border of the  $r$ - $k$  triangle is typically more diffuse, marking the highest  $k$ -values which, if combined with the corresponding  $r$ , will not result in a predicted biomass exceeding the upper prior biomass ranges. Because a large population can support a wide range of modest catch patterns even with low or medium productivity, more viable  $r$ - $k$  pairs are found in the upper left *low*-*high*- $k$  corner of the log  $r$ - $k$  space. In contrast, while a small population may be able to support high catches if it has high productivity, such catches will take a large proportion of the population resulting in strong interannual fluctuations which are prone to falling outside the theoretical and prior biomass ranges. As a result, few or no viable  $r$ - $k$  pairs are found in the lower right *high*-*low*- $k$  corner of the log  $r$ - $k$  space. As a general rule, CMSY will find more viable  $r$ - $k$  pairs in stocks where catches take a small fraction of available biomass, and vice versa. In summary, the CMSY triangle is the result of the Monte Carlo filtering process within a fixed  $r$ - $k$  space and with hard prior bounds for biomass. The tip of the triangle typically transverses the expected ellipsoid cloud of viable  $r$ - $k$  pairs found by BSM from catch and abundance data. The beauty of CMSY is that it finds this area without knowledge of abundance, albeit with a non-representative distribution. Overcoming the problems created by this triangular rather than ellipsoid distribution is the main

achievement of CMSY compared with the Catch-MSY method of Martell and Froese (2013).


### Performance of CMSY

Given the limited requirement of input data consisting of catch, qualitative resilience and qualitative stock status, the predictions of CMSY are surprisingly accurate when validated with 'true' values of simulated stocks or evaluated against BSM estimates for real stocks. While not every fluctuation of simulated or observed stock abundance is traced, the overall patterns of stock development and exploitation are usually reproduced, as shown in Fig. 4 for North Atlantic swordfish (*Xiphias gladius*, Xiphiidae, Swordfish\_NA) and Arctic cod (cod-arct). A preliminary analysis suggests that CMSY will underestimate  $MSY$  and  $k$  if landings instead of catch data are used and discards are substantial. However, even with landings


data, the estimates of  $r$  and relative biomass  $B/k$  seem to correctly reflect the productivity and the status of the stock (ICES 2014).

### Using CMSY for management of data-limited stocks

The predictions of CMSY can be presented in a format useful for stock assessment and management of data-limited stocks (Appendix IV in Data S1). In the example of Baltic brill (*Scophthalmus rhombus*, Scophthalmidae, bl-2232) (Fig. 5), catches exceeded  $MSY$  in 1995 and from 2008 to 2010, but exploitation was below the  $MSY$ -level in recent years. Highly variable  $CPUE$  data are available from 2001 onward. Both CMSY and BSM predict biomass between half  $B_{msy}$  and  $B_{msy}$  in recent years. This information is summarized in a stock status graph, showing the development of the stock from the high-exploitation-low-biomass



**Figure 5** Summary of information relevant for management of Baltic brill (bl-2232). The horizontal dashed line (a) indicates  $MSY$ , and the dotted line indicates the lower confidence limit of  $MSY$ . The bold curve in (b) is the biomass predicted by CMSY, with confidence limits (dotted curves). The normal curve indicates  $CPUE$  scaled by the catchability coefficient estimated by BSM. The horizontal dashed line indicates  $B_{msy}$  and the dotted line indicates half of  $B_{msy}$ . Panel (c) shows catch over biomass predicted by CMSY (bold curve) and catch over  $CPUE$  (normal curve) scaled by catchability  $q$  estimated by BSM. The dashed horizontal line indicates exploitation compatible with  $MSY$ . Panel (d) shows the development of biomass and exploitation relative to  $B_{msy}$  (horizontal dashed line) and  $u_{msy}$  (vertical dashed line), respectively. The horizontal dotted line indicates the biomass ( $0.5 B_{msy}$ ) below which recruitment may be impaired, and the rhomb indicates the final year in the time series. The bold curve refers to CMSY and the normal curve to BSM estimates. [CMSY\_46eFig 5-6\_2.R].



**Figure 6** Summary of information relevant for management of blond ray in the Northeast Atlantic (ICES area IXa). See Fig. 5 for general explanation of graphs (a–d). Note that CMSY curves are in bold. CPUE and catch over CPUE are plotted against their own axes in (b) relative biomass and (c) exploitation rate because the time series was too short for BSM analysis. [CMSY\_46eFig 5\_6\_2.R].

danger zone in the lower right quadrant of the graph towards the recovery zone in the upper half of the lower left quadrant. If the goal is stock recovery with better future yields, the management advice from this analysis is straightforward: maintain catches at their current low level until both CMSY and BSM predict biomass above  $B_{msy}$  for two to three years in a row and then increase catches to the lower confidence limit of MSY.

Management advice is less clear for the data-limited stock of blond ray in ICES Division IXa (*Raja brachyura*, Rajidae, rjh-pore) (Fig. 6), where CPUE data are only available from 2008 to 2014, too few for BSM analysis. CMSY predicts that catches were near MSY until 2005 and dropped to below half of MSY thereafter. Biomass recovers towards  $B_{msy}$  in 2014; however, there is a wide margin of uncertainty around that prediction. CPUE data show little change in biomass from 2008 to 2013 but confirm a drop in exploitation rate. Precautionary management may restrict catches at current levels until additional CPUE data allow for a BSM analysis and confirm a recovery to  $B_{msy}$ . At that point, the lower 95% confidence limit of MSY can serve as guidance for allowed catches.

If no CPUE data are available, other indicators can be used to confirm the predictions of CMSY analysis before they are used to inform management. For example, mean length in the catch relative to length at first maturity and relative to length at maximum cohort biomass can be used to derive independent evidence of stock status (ICES 2014, 2015; Jardim *et al.* 2014; Froese *et al.* 2015).

Given the renewed interest in the MSY concept, it may be worthwhile to repeat the following warning of the Food and Agriculture Organization of the United Nations (FAO), given at an expert consultation on the regulation of fishing effort in Rome, 17–26 January 1983: ‘Attempts to tune a system for attainment of maximum output (MSY) will lead to oscillation, unpredictability and, because of the inertia of the socio-economic system, eventually to crashes (whether reversible or not). A lower level of output is safer and more predictable’ (Caddy 1984). This warning is confirmed by the recent exploitation history of the 128 fully assessed stocks examined in this study: maximum catches had exceeded MSY in 92% of the stocks, resulting in recent biomass below the level that can produce MSY in 58% and potentially reduced



recruitment ( $B/k < 0.25$ ) in 20% of these stocks. Four stocks (3%) were severely depleted ( $B/k < 0.1$ ). In contrast, all of the 10 stocks where exploitation was kept below  $MSY$  had recent biomass levels above the one that can produce  $MSY$ , as predicted by the expert consultation in 1983 (Caddy 1984).

## Conclusions

This study presents a Monte Carlo method (CMSY) for estimating fisheries reference points from catch, resilience and qualitative stock status in data-limited stocks. It also presents a new Bayesian state-space implementation of the Schaefer production model (BSM), fitted to catch and biomass or  $CPUE$ . Both methods consider reduced recruitment and thus reduced productivity at low stock sizes and gave good predictions of  $r$ ,  $k$  and  $MSY$  when validated against simulated data. CMSY provides, in addition, reasonable predictions of relative biomass and exploitation rate when compared with 'true' simulated data. Both models were also evaluated against 128 real stocks, where estimates of biomass were available from full stock assessments. BSM estimates of  $r$ ,  $k$  and  $MSY$  were used as benchmarks for the respective CMSY estimates. These estimates were not significantly different in 76% of the stocks. A similar test against 28 data-limited stocks, where  $CPUE$  instead of biomass was available, shows that BSM and CMSY estimates were not significantly different in 89% of the stocks. Examples for using CMSY in the management of data-limited stocks are given.

## Acknowledgements

Gianpaolo Coro acknowledges the Pew Charitable Trusts for having supported his participation to the ICES WKLife 4 meeting, 27–31 October 2014, in Lisbon, Portugal. Rainer Froese acknowledges support by the European Union's Seventh Framework Programme (FP7/2007–2013) under grant agreement 244706/ECO-KNOWS project and by the Lenfest Ocean Program at The Pew Charitable Trusts under contract ID 00002841. Nazli Demirel acknowledges support from The Scientific and Technological Research Council of Turkey (TUBITAK). Kristin Kleisner acknowledges support through the National Oceanographic and Atmospheric Administration's Northeast Fisheries Science

Center and The Nature Conservancy with a grant from the Gordon and Betty Moore Foundation. This is FIN contribution number 200.

## References

- Barrowman, N.J. and Myers, R.A. (2000) Still more spawner–recruitment curves: the hockey stick and its generalizations. *Canadian Journal of Fisheries and Aquatic Sciences* **57**, 665–676.
- Beddington, J.R. and Cooke, J. (1983) *The potential yield of previously unexploited stocks*. FAO Fisheries Technical Paper No. 242, 63 pp.
- Beverton, R.J.H. and Holt, S.J. (1957) *On the Dynamics of Exploited Fish Populations*. Great Britain Ministry of Agriculture, Fisheries and Food, London.
- Caddy, J.F. (1984) *An Alternative to Equilibrium Theory for Management of Fisheries*. In FAO (1984) Papers presented at the Expert Consultation on the regulation of fishing effort (fishing mortality). Rome, 17–26 January 1983. A preparatory meeting for the FAO World Conference on fisheries management and development. FAO Fisheries Report No. 289 Suppl. 2, 214 p.
- Carruthers, T.R., Punt, A.E., Walters, C.J. et al. (2014) Evaluating methods for setting catch limits in data-limited fisheries. *Fisheries Research* **153**, 48–68.
- CFP (2013) Regulation (EU) No 1380/2013 of the European Parliament and of the Council of 11 December 2013 on the Common Fisheries Policy, amending Council Regulations (EC) No 1954/2003 and (EC) No 1224/2009 and repealing Council Regulations (EC) No 2371/2002 and (EC) No 639/2004 and Council Decision 2004/585/EC. Official Journal of the European Union L 354, 22–61. Available at: <http://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:2013:354:0022:0061:EN:PDF> (accessed 19 December 2014)
- Cope, J. and Punt, A.E. (2009) Length-based reference points for data-limited situations: applications and restrictions. *Marine and Coastal Fisheries: Dynamics, Management, and Ecosystem Science* **1**, 169–186.
- DAFF (2007) *Commonwealth Fisheries Harvest Strategy: Policy and Guidelines*. Australian Government, Department of Agriculture, Fisheries and Forestry, 55 p. Available at: <http://www.agriculture.gov.au/SiteCollectionDocuments/fisheries/domestic/hsp.pdf> (accessed 19 December 2014)
- Dick, E.J. and MacCall, A.D. (2011) Depletion-based stock reduction analysis: a catch-based method for determining sustainable yields for data-poor fish stocks. *Fisheries Research* **110**, 331–341.
- Fox, W.W. (1970) An exponential yield model for optimizing exploited fish populations. *Transaction of American Fisheries Society* **99**, 80–88.
- Froese, R. and Pauly, D. eds (2015) *FishBase. World Wide Web electronic publication*. [www.fishbase.org](http://www.fishbase.org), version

- (10/2015), accessed at [www.fishbase.org](http://www.fishbase.org) in November/December 2015.
- Froese, R., Palomares, M.L.D. and Pauly, D. (2000) Estimation of life-history key facts. In: *FishBase 2000: Concepts, Design and Data Sources* (eds R. Froese and D. Pauly), ICLARM, Philippines, p. 167–175; 344 p.
- Froese, R., Branch, T.A., Proelß, A., Quaas, M., Sainsbury, K. and Zimmermann, C. (2011) Generic harvest control rules for European fisheries. *Fish and Fisheries* **12**, 340–351.
- Froese, R., Zeller, D., Kleisner, K. and Pauly, D. (2012) What catch data can tell us about the status of global fisheries. *Marine Biology* **159**, 1283–1292.
- Froese, R., Demirel, N. and Sampang, A. (2015) An overall indicator for the good environmental status of marine waters based on commercially exploited species. *Marine Policy* **51**, 230–237.
- Froese, R., Coro, G., Kleisner, K. and Demirel, N. (2016a) Revisiting safe biological limits in fisheries. *Fish and Fisheries* **17**, 193–209.
- Froese, R., Winker, H., Gascuel, D., Sumalia, U.R. and Pauly, D. (2016b) Minimizing the impact of fishing. *Fish and Fisheries* **17**, 785–802.
- Gelman, A., Carlin, J., Stern, H. and Rubin, D. (1995) *Bayesian Data Analysis*. Chapman and Hall, New York.
- Haddon, M., Klaer, N., Smith, D.C., Dichmont, C.D. and Smith, A.D.M. (2012) *Technical Reviews for the Commonwealth Harvest Strategy Policy*. FRDC 2012/225. CSIRO, Hobart. 69 p.
- ICES (2014) *Report of the Workshop on the Development of Quantitative Assessment Methodologies Based on Life-history Traits, Exploitation Characteristics, and Other Relevant Parameters for Data-limited Stocks (WKLIFE IV)*, 27–31 October 2014, Lisbon, Portugal. ICES CM 2014/ACOM:54, 243 pp.
- ICES (2015) *Report of the Fifth Workshop on the Development of Quantitative Assessment Methodologies Based on Life-history Traits, Exploitation Characteristics and Other Relevant Parameters for Data-limited Stocks (WKLIFE V)*, 5–9 October 2015, Lisbon, Portugal. ICES CM 2015/ACOM:56. 157 pp.
- Jardim, E., Azevedo, M. and Brites, N.M. (2014) Harvest control rules for data limited stocks using length-based reference points and survey biomass indices. *Fisheries Research* **171**, 12–19.
- Jensen, A.L. (1996) Beverton and Holt life history invariants result from optimal trade-off of reproduction and survival. *Canadian Journal of Fisheries and Aquatic Sciences* **53**, 820–822.
- Kawasaki, T. (1980) Fundamental relations among the selections of life history in the marine teleosts. *Bulletin of the Japanese Society of Scientific Fisheries* **46**, 289–293.
- Kenchington, T.J. (2014) Natural mortality estimators for information-limited fisheries. *Fish and Fisheries* **15**, 533–562.
- Knezevic, A. (2008) *Overlapping Confidence Intervals and Statistical Significance*. StatNews: Cornell University Statistical Consulting Unit, 73. Available at: <https://www.cscu.cornell.edu/news/statnews/stnews73.pdf> (accessed 9 May 2016).
- Kruschke, J.K. (2011) *Doing Bayesian data analysis: a tutorial with R and BUGS*. Academic Press, Burlington, USA. 653 p.
- Lee, H.H., Maunder, M.N., Piner, K.R. and Methot, R.D. (2011) Estimating natural mortality within a fisheries stock assessment model: an evaluation using simulation analysis based on twelve stock assessments. *Fisheries Research* **109**, 89–94.
- Lee, H.H., Maunder, M.N., Piner, K.R. and Methot, R.D. (2012) Can steepness of the stock-recruitment relationship be estimated in fishery stock assessment models? *Fisheries Research* **125–126**, 254–261.
- Ludwig, D. and Walters, C.J. (1989) A Robust Method for Parameter Estimation from Catch and Effort Data. *Canadian Journal of Fisheries and Aquatic Sciences* **46**, 137–144.
- MacCall, A.D. (2009) Depletion-corrected average catch: a simple formula for estimating sustainable yields in data-poor situations. *ICES Journal of Marine Science* **66**, 2267–2271.
- Mangel, M., MacCall, A.D., Brodziak, J. et al. (2013) A perspective on steepness, reference points, and stock assessment. *Canadian Journal of Fisheries and Aquatic Sciences* **70**, 930–940.
- Martell, S. and Froese, R. (2013) A simple method for estimating MSY from catch and resilience. *Fish and Fisheries* **14**, 504–514.
- McAllister, M.K., Pikitch, E.K. and Babcock, E.A. (2001) Using demographic methods to construct Bayesian priors for the intrinsic rate of increase in the Schaefer model and implications for stock rebuilding. *Canadian Journal of Fisheries and Aquatic Sciences* **58**, 1871–1890.
- Meyer, R. and Millar, R.B. (1999) BUGS in Bayesian stock assessment. *Canadian Journal of Fisheries and Aquatic Sciences* **56**, 1078–1086.
- MFNZ (2008) *Harvest Strategy Standard for New Zealand Fisheries*. Ministry of Fisheries, Wellington, New Zealand, 27 pp. Available at: <http://www.fish.govt.nz> (accessed 19 December 2014).
- Millar, R.B. and Meyer, R. (1999) *Nonlinear State-space Modeling of Fisheries Biomass Dynamics using Metropolis-Hastings within Gibbs Sampling*. Technical Report STAT9901. Department of Statistics, University of Auckland, Auckland, New Zealand, 33 p.
- MSA (2007) *Magnuson-Stevens Fishery Conservation and Management Act, Public Law 94–265*. As amended by the Magnuson-Stevens Fishery Conservation and Management Reauthorization Act (P.L. 109–479), Available at: [http://www.nmfs.noaa.gov/msa2005/docs/MSA\\_amended\\_msa%20\\_20070112\\_FINAL.pdf](http://www.nmfs.noaa.gov/msa2005/docs/MSA_amended_msa%20_20070112_FINAL.pdf) (accessed 19 December 2014).

- Myers, R.A., Rosenberg, A.A., Mace, P.M., Barrowman, N. and Restrepo, V.R. (1994) In search for thresholds of recruitment overfishing. *Canadian Journal of Fisheries and Aquatic Sciences* **51**, 191–205.
- Myers, R.A., Bowen, K.G. and Barrowman, N.J. (1999) Maximum reproductive rate of fish at low population sizes. *Canadian Journal of Fisheries and Aquatic Sciences* **56**, 2404–2419.
- Ono, K., Punt, A.E. and Rivot, E. (2012) Model performance analysis for Bayesian biomass dynamic models using bias, precision and reliability metrics. *Fisheries Research* **125–126**, 173–183.
- Owen, A.B. (2013) *Monte Carlo Theory, Methods and Examples*. Online Stanford University publication. Available at <http://statweb.stanford.edu/~owen/mc/> (accessed 14 June 2016).
- Pauly, D. (1980) On the interrelationships between natural mortality, growth parameters, and mean environmental temperature in 175 fish stocks. *Journal du Conseil International pour l'Exploration de la Mer* **39**, 175–192.
- Pella, J.J. and Tomlinson, P.K. (1969) A generalized stock production model. *Bulletin of the Inter-American-Tropical Tuna Commission* **13**, 419–496.
- Plummer, M. (2003) JAGS: a program for analysis of Bayesian graphical models using Gibbs sampling. In: *Proceedings of the 3rd International Workshop on Distributed Statistical Computing (DSC 2003)* March 20–22, Vienna, 2003. (eds K. Hornik, F. Leisch and A. Zeileis), Vienna Technical University, Vienna pp. 20–22.
- Punt, A.E. (2003) Extending production models to include process error in the population dynamics. *Canadian Journal of Fisheries and Aquatic Sciences* **60**, 1217–1228.
- Punt, A.E. and Hilborn, R. (1997) Fisheries stock assessment and decision analysis: the Bayesian approach. *Reviews in Fish Biology and Fisheries* **7**, 35–63.
- Punt, A.E., Smith, A.D.M., Smith, D.C., Tuck, G.N. and Klaer, N.L. (2013) Selecting relative abundance proxies for BMSY and BMEY. *ICES Journal of Marine Science* **71**, 469–483.
- Ricker, W.E. (1975) *Computation and Interpretation of Biological Statistics of fish Populations*. Bulletin of the Fisheries Research Board of Canada 191, Ottawa, Canada, 382 pp.
- Rosenberg, A.A., Fogarty, M.J., Cooper, A. B. et al. (2014) *Developing New Approaches to Global Stock Status Assessment and Fishery Production Potential of the Seas*, FAO Fisheries and Aquaculture Circular No. 1086 FIR/C1086, FAO, Rome.
- Schaefer, M. (1954) Some aspects of the dynamics of populations important to the management of the commercial marine fisheries. *Bulletin of the Inter-American-Tropical Tuna Commission* **1**, 27–56.
- Schnute, J.T. and Richards, L.J. (2002) Surplus production models. In: *Handbook of Fish Biology and Fisheries*, Vol. **2**. (eds P.J.B. Hart and J.D. Reynolds), Blackwell Publishing, Oxford, UK, pp. 105–126.
- Smith, R.W. (1995) *Visual Hypothesis Testing with Confidence Intervals*. EcoAnalysis Inc., Ojai, CA, 1–6.
- Sparre, P. and Venema, S.C. (1998) *Introduction to Tropical Fish Stock Assessment*. Part 1. Manual. FAO Fisheries Technical Paper No. 306.1, Rev.2., Rome, FAO, 407 pp.
- Thorson, J.T., Branch, T.A. and Jensen, O.P. (2012) Using model-based inference to evaluate global fisheries status from landings, location, and life history data. *Canadian Journal of Fisheries and Aquatic Sciences* **69**, 645–655.
- Thorson, J.T., Minto, C., Minto-Vera, C.V., Kleisner, K.M. and Longo, C. (2013) A new role for effort dynamics in the theory of harvested populations and data-poor stock assessment. *Canadian Journal of Fisheries and Aquatic Sciences* **70**, 1829–1844.
- Thorson, J.T., Ono, K. and Munch, S.B. (2014) A Bayesian approach to identifying and compensating for model misspecification in population models. *Ecology* **95**, 329–341.
- Vasconcellos, M. and Cochrane, K. (2005) Overview of world status of data-limited fisheries: inferences from landing statistics. In: *Fisheries Assessment and Management in Data-limited Situations*. (eds G.H. Kruse, V.F. Gallucci, D.E. Hay, R.I. Perry, R.M. Peterman, T.C. Shirley, P.D. Spencer, B. Wilson and D. Woodby), Alaska Sea Grant Programme, Fairbanks, AK, USA, pp. 1–20.
- Zhou, S., Yin, S., Thorson, J.T., Smith, A.D.M. and Fuller, M. (2012) Linking fishing mortality reference points to life history traits: an empirical study. *Canadian Journal of Fisheries and Aquatic Sciences* **69**, 1292–1301.

## Supporting Information

Additional Supporting Information may be found in the online version of this article:

**Data S1.** Supplementing information consists of a detailed representation of the results of all stock analyses, including links to data and R-code.