



18th World Congress of the International Association of Surgeons, Gastroenterologists and Oncologists



October 08 - 11, 2008, İstanbul / Türkiye
Lütfi Kırdar Convention and Exhibition Center

Congress President
Dr. Yusuf Bayraktar

Honorary Presidents
Dr. Faruk Memik
Dr. Masatoshi Makuuchi
Dr. Nick J. Lygidakis

Organizing Committee

Dr. İsmail Arslan
Dr. Semih Baskan
Dr. Adil Baykan
Dr. Yılmaz Çakaloğlu
Dr. Mustafa Keçer
Dr. Bülent Sivri
Dr. Hakan Şentürk
Dr. İlkay Şimşek
Dr. Nurdan Tözün
Dr. Serhat Ünal
Dr. Şuayib Yalçın
Dr. Abdullah Zorluoğlu

Scientific Secretariat



Hacettepe University School of Medicine
Department of Gastroenterology

Phone : +90 312 305 14 90
+90 312 442 94 28
Fax : +90 312 442 94 29
E-mail : bayrak@hacettepe.edu.tr
bayraktar_yusuf@hotmail.com

Organization Secretariat



Serenas Tourism Congress
Organization Services Ltd. Co.

Turan Güneş Bulvarı 5. Cad. No:13, 06550
Yıldız, Çankaya - ANKARA, TÜRKİYE
Phone : +90 312 440 50 11
Fax : +90 312 441 45 61
E-mail : iasgo@serenas.com.tr
Url : www.serenas.com.tr

www.iasgo2008.org



A NEW MINIMALLY INVASIVE TECHNIQUE FOR THE TREATMENT OF NECROTIZING PANCREATITIS: ENDOSCOPIC RETROPERITONEAL NECROSECTOMY AND DEBRIDEMENT (ERND)**Afkaan Gök, Cemalettin Ertekin, Hakan Yanar, Recep Güloğlu, Korhan Taviloğlu***Istanbul Medical Faculty*

Objectives: Laparotomy is performed in patients with necrotizing pancreatitis, if their clinical condition is adjudged to be worsening or multiple organ failure ensues. As an alternative to laparoscopic techniques performed under general anaesthesia in the retroperitoneal space, endoscopic techniques can be used, in which local anaesthesia and sedation are sufficient to spare the patient additional surgical stress.

Patients and Methods: Between November 2006 and October 2007, 3 of 41 patients presenting with proven infected pancreatic necrosis underwent endoscopic retroperitoneal necrosectomy and debridement (ERND). A 15 mm laparoscopic trocar was inserted with CT guidance under local anaesthesia into patients who had a fluid collection in the left retroperitoneal area. Necrosectomy and drainage were performed using a double-channel gastroscope and a basket catheter. After ERND, 10mm aspirative silicone drainage tube was inserted into the peripancreatic area under endoscopic guidance.

Results: Of the 7 patients from the sample of 41 who underwent open necrosectomies and drainage, 3 died (43%). Of the 3 who had ERND, 1 patient died in post-operative intensive care two weeks after the procedure (33%). Cause of death was due to the patient's critical condition and cardiopulmonary complications which rendered patient inappropriate for general anaesthesia. All of 3 patients in the ERND group were confirmed on CT scan to be free of necrotic material in the peripancreatic area post-operatively. One patient developed pancreatic fistula for which Wirsungotomy was performed. None of the patients suffered hollow viscus or vascular damage due to ERND.

Conclusion: ERND provides an effective approach to the treatment of necrotizing pancreatitis which forgoes need for open surgery under general anaesthesia. We believe that the difference will be more apparent as number of ERND cases increase since the patient goes under decreased stress due to the avoidance of general anesthesia and open surgery.