

**第113回**  
**日本林学会**  
**大会**  
**学術講演集**

**2002**

## 目 次

国際講演会「環日本海沿岸諸国(ロシア、中国、韓国、日本)における森林資源の現状と課題」 講演要旨	
Problems of forest management in Siberia .....	V.A. Sokolov (Sukachev Inst., Russia) 1
Supply and demand of forest resources in South Korea	
.....	Jong-Cheon Choi (Kangwon Univ., South Korea) 3
Forest resources and problems in Niigata Prefecture	
.....	Masao Noomote (Niigata Pref. For. Inst., Japan) 6

### 学会賞受賞者講演要旨

#### 日本林学会賞

アメリカ国有林管理の史的展開 .....	大田伊久雄 (京大農) 9
松枯れをめぐる宿主-病原体-媒介者の相互作用 .....	富樫一巳 (広島大総合科) 11
森林作業と労働負担 .....	井上公基 (日大生物資源) 12

#### 日本林学会奨励賞

相対成長式によるスギ同齢単純林における樹高曲線の解析 .....	井上昭夫 (鳥取大農) 13
落葉広葉樹林におけるN <sub>2</sub> O生成とその空間的異質性 .....	石塚成宏 (森林総研北海道) 14

### 口頭発表・部門別

#### 造 林

セコイアオスギ ( <i>Sequoiadendron giganteum</i> Buch) の育苗III .....	奥村覚 (林学会員) 15
高速道路法面に植栽されたトウヒ属樹木の成長反応 .....	香山雅純ら (北大生物圏セ) 16
マレーシアサラワク州ニア造林試験地に植栽された7樹種の生態生理 .....	田中憲蔵ら (愛媛大農) 17
ブナ苗木の樹幹直径日変化と蒸散量の関係 .....	鍛代邦夫ら (日大生物資源) 18
熱帯林の樹冠部における水利用特性 -マレーシア ランビルヒルズ国立公園の例-	
.....	北橋善範ら (北大院農) 19
根系への硫酸酸性雨処理がモミ成木の生理、生態に及ぼす影響 .....	英賀慶彦ら (東大院農) 20
異なる水分とCO <sub>2</sub> 条件で生育したシラカンバの光合成特性と被食防衛物質	
.....	小池孝良ら (北大生物圏セ) 21
広葉樹二次林伐採地における萌芽の成長とシカによる食害の影響 .....	小倉悠ら (東大農) 22
樹幹形成過程における密度効果 -幹の細りの表現形-	
.....	千葉幸弘 (森林総研) 23
落葉広葉樹林の相対成長関係の互換性と推定精度 .....	小見山章ら (岐阜大農) 24
東吉野村におけるスギ人工林の密度管理 -間伐直後の林分のD-H関係-	
.....	高橋絵里奈ら (京大演) 25
センダンの枝の成長特性について .....	横尾謙一郎 (熊本県林研指) 26
スズタケ地上部と地下部の現存量 .....	中本真平ら (信州大農) 27
11年生ミズメ人工林の林分構造と現存量 .....	松山知恵ら (京大農) 28
ヒノキ不成績人工林におけるヒノキと侵入広葉樹の成長解析 -京都府美山町の24年生ヒノキ人工林を事例として-	
.....	鳴田俊平ら (京大農) 29
カラマツ不成績造林地林床の光環境の季節変化に対する広葉樹稚樹の応答	
.....	北岡哲ら (北大院農) 30
ヒノキ人工林におけるギャップ周辺の光環境とヒノキ天然生稚樹の動態 .....	松崎誠司ら (東農大院) 31

MuPSマーカーを用いたヤマナラシ ( <i>Populus tremula</i> ) のクローン構造解析 -クローナル木本におけるラミートとジェネットの整合性-	本間航介ら (新潟大農フ科セ)	616
韓国中部地域における天然落葉広葉樹林の土壌CO <sub>2</sub> の発生	陳鉉五ら (慶熙大)	617
天然更新樹木稚苗の根系構造 -特性解明のための指標化-	溝口岳男ら (森林総研)	618
アルカリ吸着法、通気法および密閉式循環法で測定された土壌呼吸速度の比較	大橋瑞江ら (九大院農)	619
樹皮呼吸測定装置の開発	角張嘉孝ら (静岡大農)	620
腐生性担子菌の落葉分解と水分要求	小出奏ら (東京農工大農)	621
アオダモ種子の発芽阻害要因	滝谷美香ら (北海道林試)	622

## 生理 I・II

中国半乾燥地における凝結水について	李玉壺ら (岡山大院)	623
Growth responses of several tree species to flooding stress	Zülfikar, Halide Candanら (岡山大院自然科学)	624
四日市市におけるシデコブシの葉の水分特性と水ポテンシャルの日変化	斎場勇治ら (三重大生物資源)	625
生育環境が苗木の生理特性に及ぼす影響	角園敏郎 (森林総研)	626
外生菌根菌の接種がアカマツ実生苗の生長および水分生理に及ぼす影響	三輪誠ら (埼玉県環科国セ)	627
<i>Acacia ampliceps</i> 芽生えにおける塩ストレスによるカタラーゼの発現誘導について	横田智 (森林総研)	628
<i>Morus alba</i> (カラヤマグワ) と <i>Morus bombycis</i> (ヤマグワ) の外樹皮、内樹皮成分の定量的差違について	石井洋二ら (福島県林研セ)	629
ジベレリン生合成処理がメヒルギの発根および根の成長におよぼす効果	柘田信彌ら (東農大地域環境)	630
葉内 $\alpha$ -カロチン含量の樹種間差と季節変動について	柴田勝ら (長岡高専)	631

## 育種 I・II

稀少樹種ハナノキの遺伝変異 (予報)	金指あや子ら (森林総研)	632
シデコブシ、コブシ、タムシバの形態調査とアロザイム解析	中島美幸ら (岐阜県森林研)	633
モミの母樹間の距離と種子の有胚率およびアロザイム変異	斉藤陽子ら (東大樹芸研)	634
イヌブナの遺伝的変異と集団分化	鳥居由季子ら (名大院生命農)	635
SSRマーカーによるヒノキ精英樹の識別と次代検定林の花粉親推定	森康浩ら (福岡県林技セ)	636
クロマツのマツバノタマバエ抵抗性と連鎖したRAPDマーカーのSCARマーカー化	林英司ら (林育セ)	637
ハゼノキのクローン別特性評価 -RAPDマーカーによるクローン分類と木蠟収量の年次変動-	平岡裕一郎ら (林育セ九州)	638
九州各地の抵抗性クロマツ採種園から得られた「田辺ク-54号」家系苗の花粉親とマツノサイセンチュウ抵抗性の関係	宮原文彦ら (福岡県林技セ)	639
ヒノキ採種園の交配実態に関するマイクロサテライト分析	中尾有里ら (名大院生命農)	640
スギ ( <i>Cryptomeria japonica</i> D.Don) のマイクロサテライトマーカーの開発とその連鎖地図の作成	高橋友和ら (新潟大院自然科学)	641
AFLP, CAPSおよびSSRマーカーによるヒノキ連鎖地図	松本麻子ら (森林総研)	642

## Growth Responses of Several Tree Species to Flooding Stress

Zulfikar, Halide Candan • Sakamoto, Keiji (Okayama University, Graduate School of Natural Science and Technology) and Yoshikawa, Ken (Okayama University, Faculty of Agriculture)

## Introduction

The relationship between water and plant productivity is extremely complex. Plants exhibit broad range of adaptations to either excesses or deficits of water. Water excesses such as flooding has effects on survival and growth of plants. Adaptation capacity of plants to flooding varies greatly among plant species.

Recently, wet site planting comes into view more necessarily because of the predicted global warming and sea level rise, erosion control around water resources and dam sites. Additionally, increase of common interest in landscaping along water-zone areas for recreation purposes is also an important factor for wet site planting. This reveals the necessity of more data on tree responses to flooding stress.

The present study reports the growth responses and resistant of saplings of *Alnus hirsuta* Trucz., *Populus alba* L. and *Taxodium distichum* L. tree species to flooding stress during the growing season.

## Material and Methods

Saplings of *A. hirsuta* (53 cm – 94 cm in height), *P. alba* (86.5 cm – 132 cm in height) and *T. distichum* (48 cm – 70 cm in height) were used for the experiment. Twenty-four saplings of each species were transplanted into plastic pots in April 1999. Experiment was conducted in the green house condition and the saplings were exposed to the flooding stress in the plastic containers with different water levels. The saplings of each species were divided into four groups as follows;

1. Six saplings were unflooded during the experiment (UF). These saplings were watered every day or every two days.
2. Six saplings were flooded to the soil surface with stagnant water in a plastic container (F<sub>0</sub>).
3. Six saplings were flooded to the height of 20 cm above the soil surface with stagnant water (F<sub>20</sub>).
4. Six saplings were flooded to the soil surface with air-involved water in a plastic container (F<sub>air</sub>).

Heights, diameters, leaf numbers and branch lengths were measured during the experiment. The heights, and stem base diameters at the height of 3 cm and 20 cm above soil surface were measured for all the saplings before flooding treatment in April 1999. They were remeasured in September and December 1999. The length of all branches of each sapling was measured once every two weeks from April to September 1999. The number of leaves of saplings was counted once every two weeks from April to September 1999.

## Results and Discussion

All the saplings of *P. alba* and *T. distichum* survived but five saplings of *A. hirsuta* were dead under the flooding conditions. Four saplings of *A. hirsuta* were dead within 3 weeks after beginning of the experiment. Three of them were in the treatment group of F<sub>0</sub>, and the other one was in the treatment group of F<sub>air</sub>. One other *A. hirsuta* sapling was dead on the 5<sup>th</sup> week of the experiment in the F<sub>20</sub> treatment group.

Flooding induced reduction in height growth in all treatment groups of the *A. hirsuta* saplings. On the other

hand, in the *P. alba* and *T. distichum* saplings, there were no significant differences between treatment groups and control group.

Flooding caused reduction in diameter increment at 3 cm height in the F<sub>0</sub> and F<sub>20</sub> treatment groups of the *A. hirsuta* saplings and F<sub>20</sub> treatment group of *P. alba* saplings. On the other hand, diameter increment at 3 cm height was greater for *T. distichum* saplings in the flooded saplings than unflooded saplings, although the difference was significant only F<sub>20</sub> treatment group. *P. alba* and *T. distichum* saplings exhibited no treatment differences in diameter increment at 20 cm height, but flooded *A. hirsuta* saplings showed significantly reduced diameter increment at 20 cm height. Ratio of diameter increment at 20 cm height to 3 cm height was significantly high in F<sub>20</sub> treatment group of *P. alba* saplings and significantly low for F<sub>0</sub> treatment group of *T. distichum* saplings (Fig. 1). It was observed that water level affected the morphological characteristics of stems in the flooding conditions. It was thought that hypertrophied lenticels were developed on submerged parts of stems and bark thickening was increased due to accelerated proliferation of phloem parenchyma cells and large amounts of intercellular space in phloem.

Flooding significantly inhibited branch length increment of *A. hirsuta*, *P. alba* and *T. distichum* saplings in the all treatment groups comparing to the control group. Decrease in branch length increment was greater in *A. hirsuta* and *P. alba* saplings than *T. distichum* saplings.

Emerging leaves before treatment exhibited no significant difference in leaves shedding between treatment groups and control group in all tree species. On the other hand, the number of emerging leaves after treatments was significantly low in treatment groups when comparing to the control group. Difference was higher in *A. hirsuta* saplings than *P. alba* and *T. distichum* saplings. The decrease in the total number of leaves was probably due to early leaf shedding and inhibition or delay of new leafing.




Fig. 1 Ratio of diameter increment at 20 cm height to 3 cm height (Turkey test  $p < 0.05$ )